



# **KOD Hot Start DNA Polymerase**

KOD Hot Start DNA Polymerase

Page 1 of 8

20 U	71086-5
200 U	71086-3
1000 U	71086-4

### **About the Kits**

### Description

KOD Hot Start DNA Polymerase is a premixed complex of the high fidelity KOD DNA Polymerase and two monoclonal antibodies that inhibit the DNA polymerase and 3' $\rightarrow$ 5' exonuclease activities at ambient temperatures (1). KOD Hot Start combines the high fidelity, fast extension speed, and outstanding processivity of KOD with the high specificity of an antibody-mediated hot start. Non-specific amplification is reduced because mispriming events during reaction set up and the initial temperature increase are avoided. In addition, primer degradation during setup at ambient temperature due to exonuclease activity is effectively inhibited. This enzyme quickly and accurately amplifies genomic and phage/plasmid DNA targets up to 12 and 20 kbp, respectively. GC-rich targets are also efficiently amplified. KOD Hot Start DNA Polymerase produces blunt-ended DNA products that are suitable for cloning with the Novagen Perfectly Blunt<sup>®</sup> and LIC Vector Kits and is compatible with site-directed mutagenesis protocols.

**Unit definition:** One unit is defined as the amount of enzyme that will catalyze the incorporation of 10 nmol of dNTP into acid insoluble form in 30 minutes at 75°C in a reaction containing 20 mM Tris-HCl (pH 7.5 at 25°C), 8 mM MgCl<sub>2</sub>, 7.5 mM DTT, 50  $\mu$ g/ml BSA, 150  $\mu$ M each of dATP, dCTP, dGTP, dTTP (a mix of unlabeled and [<sup>3</sup>H]-dTTP) and 150  $\mu$ g/ml activated calf thymus DNA.

© 2006 EMD Biosciences, Inc., an affiliate of Merck KGaA, Darmstadt, Germany. All rights reserved. Perfectly Blunt<sup>®</sup> and the Novagen<sup>®</sup> name is a registered trademark of EMD Biosciences, Inc. in the United States and in certain other jurisdictions. TWEEN<sup>®</sup> is a registered trademark of ICI Americas Inc. *PfuTurbo<sup>®</sup>* and *PfuUltra<sup>®</sup>* are registered trademarks of Stratagene. KOD Polymerases are manufactured by TOYOBO and distributed by EMD Biosciences, Inc., Novagen. KOD XL DNA Polymerase is licensed under U.S. Patent No. 5,436,149 owned by Takara Shuzo, Co., Ltd.

Use of this product is covered by one or more of the following US patents and corresponding patent claims outside the US: 5,079,352, 5,789,224, 5,618,711, 6,127,155, 5,677,152, 5,773,258, and claims outside the US corresponding to US Patent No. 4,889,818. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patents claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim (such as the patented 5' Nuclease Process claims in US Patent Nos. 5,210,015 and 5,487,972) and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other comical consideration, is conveyed expressly, by implication, or by estoppel. This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained by contacting the Direction of Licensing, Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404, USA.

USA and Canada Tel (800) 526-7319 novatech@novagen.com Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk All Other Countries Contact Your Local Distributor www.novagen.com novatech@novagen.com

A Brand of EMD Biosciences, Inc., an Affiliate of Merck KGaA, Darmstadt, Germany www.novagen.com

FOR RESEARCH USE ONLY. NOT FOR HUMAN OR DIAGNOSTIC USE.

#### Polymerase fidelity comparison



#### Mutation frequency comparison: KOD Hot Start, *PfuTurbo<sup>®</sup>*, *PfuUltra<sup>®</sup>*, and *Taq*

The fidelity of replication was measured as the mutation frequency in PCR products using a modified  $rpsL^+$  fidelity assay (2, 3).

#### Polymerase rate comparison

Enzyme	KOD DNA Polymerase	<i>Pfu</i> DNA Polymerase	<i>Taq</i> DNA Polymerase
Species	Thermocccus kodakaraensis	Pyrococcus furiosus	Thermus aquaticus YT-1
Elongation rate (bases/second)	106–138	25	61
Processivity* (nucleotide bases)	> 300	< 20	not determined

\* Processivity is defined as the number of nucleotides that can be extended in one catalytic reaction by one DNA polymerase molecule.

#### **Components**

20 U or 200 U or 5 × 200 U KOD Hot Start DNA Polymerase (1 U/µl in 50 mM Tris-HCl, 1mM DTT, 0.1 mM EDTA, 50% glycerol, 0.001% Nonidet P-40, 0.001% Tween<sup>®</sup>-20, pH 8.0)

10X PCR Buffer for KOD Hot Start DNA Polymerase

- 1.2 ml or  $5 \times 1.2$  ml
- 1 ml or  $5 \times 1$  ml 25 mM MgSO<sub>4</sub>
- 1 ml or  $5 \times 1$  ml dNTPs (2 mM each)

### Storage

Store all components at  $-20^{\circ}$ C.

### **KOD Hot Start DNA Polymerase Protocol**

KOD Hot Start DNA Polymerase and buffer are a unique PCR system. The following procedure is designed for use with the components provided in the KOD Hot Start DNA polymerase kit. Using reaction components or protocols designed for any other DNA polymerase may result in poor amplification. Reaction conditions listed below will provide satisfactory amplification for most primer/template combinations. Guidelines and troubleshooting sections provide details for optimizing reaction conditions.

Examples of amplification from human genomic DNA and plasmid DNA can be found in the Appendix on page 7.

Component	Volume	Final Concentration	
10X Buffer for KOD Hot Start DNA Polymerase	5 µl	1X	
25 mM MgSO <sub>4</sub> <sup>a</sup>	3 µl	1.5 mM	
dNTPs (2 mM each)	5 µl	0.2 mM (each)	
PCR Grade Water	Xμl		
Sense (5') Primer (10 µM)	1.5 µl	0.3 µM	
Anti-Sense (3') Primer (10 µM)	1.5 µl	0.3 μΜ	
Template DNA <sup>b</sup>	Υµl		
KOD Hot Start DNA Polymerase (1 U/µl)	1 µl	0.02U/µl	
Total reaction volume	50 µl		

### **Standard reaction setup**

<sup>a</sup> To optimize for targets greater than 2kb, final Mg<sup>2+</sup> concentration may be adjusted to between 1.5 and 2.25 mM.

<sup>b</sup> See Template DNA section on page 4.

### **Cycling conditions**

#### **Temperature and time**

The following table allows for primer extension that occurs during temperature ramping between steps.

	Target size			
Step	< 500 bp	500–1000 bp	1000–3000 bp	> 3000 bp
1. Polymerase activation	95°C for 2 min	95°C for 2 min	95°C for 2 min	95°C for 2 min
2. Denature	95°C for 20 s	95°C for 20 s	95°C for 20 s	95°C for 20 s
3. Annealing	Lowest Primer Tm°C for 10 s			
4. Extension	70°C for 10 s/kb	70°C for 15 s/kb	70°C for 20 s/kb	70°C for 25 s/kb
Repeat steps 2–4	20-40 cycles. For more information see "Cycle number" below			

#### **Cycle number**

The number of cycles (steps 2 through 4 in the above table) required to generate a PCR product will depend on the source and amount of starting template in the reaction, as well as the efficiency of the PCR. In general, 20–40 cycles will be adequate for a wide range of templates. It is common to use fewer cycles when amplifying targets from plasmids (i.e., subcloning) where a high number of copies of template is easily attained, as this reduces the chance of amplifying a mutation. A higher number of cycles (e.g., 40) may be necessary when amplifying from genomic DNA since the target sequence will be in low abundance.

USA and Canada Tel (800) 526-7319 novatech@novagen.com Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk

### Additional Guidelines

### **Primers**

Primer design is critical for successful PCR amplification. Because KOD Hot Start DNA polymerase exhibits strong  $3' \rightarrow 5'$  exonuclease activity after thermo activations, primers should be at least 21 bases of 3' end complementary to the target sequence. G/C content of the primers should be 40-60%. Primer melting temperature (T<sub>m</sub>) is defined as the temperature at which one half of the DNA duplex will dissociate to become single stranded. Some primer molecules will anneal as the temperature approaches the T<sub>m</sub> of a primer, as a result PCR amplifications are usually successful over a range of annealing temperatures. Primer pairs with similar T<sub>m</sub> values usually result in better amplifications because annealing and extension are better synchronized. If melting temperatures of a primer pair differ by more than 5°C, increasing the length of the lower-T<sub>m</sub> primer will reduce the difference.

There are several methods for determining the  $T_m$  of a primer. The nearest-neighbor method (4) using 50 mM monovalent salt is one method for  $T_m$  prediction. Unlike other methods, the nearest-neighbor method takes into account the primer sequence and other variables such as salt and DNA concentration. The  $T_m$  can also be calculated with the % GC method (5). The most general method of calculating the  $T_m$  is based on the number of adenine (A), thymidine (T), guanidine (G) or cytosine (C) bases where  $T_m(^\circ C) = 2(N_A + N_T) + 4(N_G + N_C)$ .

Primer  $T_m$  values reported by manufacturers may vary by 5 to 10°C depending on the calculation method used. In addition, the exact  $T_m$  for a given primer in a reaction may be affected by DNA concentrations (primer and template), mono and divalent ion concentrations, dNTP concentration, presence of denaturants (e.g., DMSO), and nucleotide modifications. Therefore, an optimal primer annealing temperature should be determined empirically.

When receiving oligonucleotides from the manufacturer, prepare primer stocks at 100 pmol/ $\mu$ l (100  $\mu$ M) in TE and store them at -20°C. To set up KOD reactions, dilute enough of each primer stock 10-fold (10 $\mu$ M) to add 1.5  $\mu$ l per reaction.

#### **Template DNA**

The optimal amount of starting template may vary depending on the template quality. In general the suggested amount of template DNA for amplification is 10 ng phage DNA, 10 ng plasmid DNA, 100 ng genomic DNA, or 2  $\mu$ l of a reverse transcription reaction. Using too much template in the PCR reaction can result in failed reactions since template denaturation is concentration dependant. At high concentrations of DNA, denaturation is less efficient.

#### **Plasmid templates**

For subcloning, amplify from 10 ng of plasmid template and reduce the number of cycles to 20-25.

#### **GC-rich templates**

The addition of DMS0 to 2-10% final concentration may decrease template secondary structure and increase yield. Final DMSO concentrations of less than 5% v/v have no effect on fidelity (6, 7). The effect of DMSO above 5% v/v on enzyme fidelity has not yet been determined.

#### Unpurified templates

Crude cell lysates, PCR products, plaques, and colonies can serve as template for PCR. Limit the volume of unpurified templates to reduce inhibition of the reaction.

#### Long target DNA

Amplification of targets longer than 3000 bp can be improved by increasing the concentration of MgSO<sub>4</sub>. Adjusting the final MgSO<sub>4</sub> concentration from 1.5 to 2.25mM in 0.25 mM increments should be tried when suboptimal results are obtained for targets over 3000 bp. Also, the addition of DMSO to 2–10% v/v final concentration may reduce secondary structure of the template DNA and increase yield.

Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk

#### **Reaction components**

High volumes of primer and template DNA suspended in Tris-EDTA (TE) will chelate free  $Mg^{2+}$  in the reaction and may affect enzyme performance. If the combined volume of primer and template in TE in the reaction exceeds 5 µl, then adjustment of the  $MgSO_4$  concentration may be necessary. Increase the  $MgSO_4$  in the reaction to compensate for the EDTA. Each molecule of EDTA will chelate one molecule of  $Mg^{2+}$ , so increasing the  $MgSO_4$  by 0.25µl (0.125mM) for every 6.3µl of TE will compensate for the  $Mg^{2+}$  chelated by the EDTA.

### **Extension temperature and time**

Extension at 70°C is recommended since a good balance of polymerization speed and accuracy is obtained. KOD Hot Start DNA Polymerase exhibits optimal proof reading activity at 68°C and optimal polymerization activity at 74°C. If using an extension temperature near 74°C, shortening the extension time 5 s/kbp may give better amplification. When using an extension temperature near 68° C, increasing the extension time by 5 s/kbp may give better results.

#### **Two-step PCR**

In two-step PCR, annealing and extension can be carried out at the same temperature. Primers for two-step cycling programs should be designed with high  $T_m$  values (> 65°C) to ensure proper annealing and extension at the same temperature. Initially try an annealing/extension temperature equal to the lowest  $T_m$  of the primer pair. Since polymerase speed is slower at 68°C, increase the annealing/extension time by 5 s/kbp during two-step cycling.

### **Optimization**

When optimizing PCR reactions, it is best to change only one parameter at a time. The use of DMSO at 5% v/v final often improves a suboptimal PCR.

## Troubleshooting

Symptom	Possible cause	Solution
No PCR product	Extension time is too long	Lower extension time to 15 s/kpb
	Too much secondary structure in template DNA	Add DMSO to a final concentration of 5–10% v/v
	PCR primers are not long enough	Use primers longer than 21 bases
	Annealing temperature is too high	Lower annealing temperature in 3°C decrements
Low yield	High GC content	Add DMSO to a final concentration of 5–10% v/v.
	Suboptimal PCR conditions	Increase final MgSO4 concentration in 0.25 mM increments.
	Long target DNA	Increase final MgSO $_4$ concentration in 0.25 mM increments.
Smearing	Too much template DNA	Reduce the amount of template DNA
Smearing below target size	Extension times are too short	Increase extension time 5 s/kbp
	MgSO <sub>4</sub> concentration too low	Increase final MgSO <sub>4</sub> concentration 0.25 mM increments
Smearing above target size	Extension times too long	Reduce extension time 5 s/kbp
	MgSO₄ concentration too high	Decrease final MgSO <sub>4</sub> concentration in 0.25 mM increments
Primer dimers	Primers are complementary to each other	Design primers that are not self-complementary or complementary to each other
	Primer concentration is too high	Reduce primer concentration
	Annealing temperature too low	Raise annealing temperature

# **Application references**

This section lists selected references for applications with KOD Hot Start DNA Polymerase. Please visit <u>www.novagen.com/KOD</u> for the latest information.

Application	Reference
Colony-direct PCR with Gram-positive bacteria	Tsuchizaki, N. and Hotta, K. (2003) <i>inNovations</i> <b>17</b> , 9–11.
Elongation PCR	Gao, X., Yo, P., Keith, A., Ragan, T. J., and Harris, T. K. (2003) <i>Nucleic Acids Res.</i> <b>31</b> , e143.
Gene cloning Schilling, O., Spath, B., Kostelecky, B., Marchfelder, A., Meyer-Klaucke, W., and Vogel, A. (2005) <i>J. Chem.</i> <b>280</b> , 17857–17862.	
	Williams, M. E., Burton, B., Urrutia, A., Shcherbatko, A., Chavez-Noriega, L. E., Cohen, C.J., and Aiyar, J. (2005) <i>J. Biol. Chem.</i> <b>280</b> , 1257–1263.
	Miyazato, T., Toma, C., Nakasone, N., Yamamoto, K., and Iwanaga, M. (2003) J. Med. Microbiol. 52, 283–288.
Gene cloning using _consensus shuffling	Binkowski, B. F., Richmond, K. E., Kaysen, J., Sussman, M. R., and Belshaw, P. J. (2005) <i>Nucleic Acids Res.</i> 33, e55.
Multiplex cDNA-PCR	Sagara, N. and Katho, M. (2000) <i>Cancer Res.</i> <b>60</b> , 5959–5962.
Multiplexed SNP genotyping	Higasa, K. and Hayashi, K., (2002) Nucleic Acids Res. 30, e11.
Mutagenesis Tabuchi, M., Tanaka, N., Nishida-Kitayama, H., Ohno, H., and Kishi, F. (2002) Mol. Biol. Cell 13, 4371-	
	Matsumoto, N., Mitsuki, M., Tajima, K., Yokoyama, W. M., and Yamamoto, K. (2001) <i>J. Exp. Med.</i> <b>193</b> , 147–158.
PCR for PCR-Mass spectrometry based analysis	Benson, L. M., Null, A. P., and Muddiman, D. C. (2003) J. Am. Soc. Mass. Spectrom. 14, 601–604.
PCR for sequence analysis	Okamoto, T., Yoshiyama, H., Nakazawa, T., Park, I. D., Chang, M. W., Yanai, H., Okita, K., and Shirai, M. (2002) <i>J. Antimicrob. Chemother.</i> <b>50</b> , 849–856.
Second strand cDNA synthesis	Hirohashi, Y., Torigoe, T., Maeda, A., Nabeta, Y., Kamiguchi, K., Sato, T., Yoda, J., Ikeda, H., Hirata, K., Yamanaka, N., and Sato, N. (2002) <i>Clin. Cancer Res.</i> <b>8</b> , 1731–1739.

USA and Canada Tel (800) 526-7319 novatech@novagen.com Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk

### References

۱.	Mizuguchi, H., Nakatsuji,	M., Fujiwara, S.,	Takagi, M. and	Imanaka, T. (	(1999) J. Biocher	m (Tokyo) <b>126</b> ,
	762–768.					

- Kitabayashi, M., Nishiya, Y., Esaka, M., Itakura, M., and Imanaka, T. (2002) *Biosci. Biotechnol.* Biochem. 66, 2194–2200.
- Fujii, S., Akiyama, M., Aoki, K., Sugaya, Y., Higuchi, K., Hiraoka, M., Miki, Y., Saitoh, N., Yoshiyama, K., Ihara, K., Seki, M., Ohtsubo, E., and Maki, H. (1999) *J. Mol. Biol.* 289, 835–850.
- Breslauer, K. J., Frank, R., Blocker, H. and Marky, L. A. (1986) *Proc. Natl. Acad. Sci.* 83, 3746–3750.
- 5. Howley, P. M., Israel, M. A., Law, M. F. and Martin, M. A. (1979) J. Biol. Chem. 254, 4876-4883.
- Cheng, S., Fockler, C., Barnes, W. M., and Higuchi, R. (1994) Proc. Natl. Acad. Sci. USA 91, 5695–5699.
- 7. Winship, P. R. (1989) Nucleic Acids Res. 17, 1266.

### **Appendix**

### **Example amplifications**

### Amplification of 335 bp fragment (CFTR exon 11) from human genomic DNA

Reaction setup

Component	Volume	Final Concentration
10X Buffer for KOD Hot Start DNA Polymerase	5 µl	1X
25 mM MgSO <sub>4</sub>	3 µl	1.5 mM
dNTPs (2 mM each)	5 µl	0.2 mM (each)
PCR Grade Water	32 µl	
Sense (5') Primer (10 µM)	1.5 µl	0.3 µM
Anti-Sense (3') Primer (10 µM)	1.5 µl	0.3 µM
Human Genomic DNA* (100 ng/µl)	1 µl	2 ng/µl
KOD Hot Start DNA Polymerase (1 U/µl)	1 µl	0.02U/µl
Total reaction volume	50 µl	

\* Human Genomic DNA (Cat. No. 69237-3) diluted in TE to 100 ng/ul

### Cycling conditions

Step	Temperature and time
1. Polymerase activation	95°C for 2 min
2. Denature	95°C for 20 s
3. Annealing	56°C for 10 s
4. Extension	70°C for 4 s
Repeat steps 2–4	30 cycles
5. Hold	4°C



1.2 % TAE agarose gel Lane 1 PCR Markers, 50 2000 bp (Cat. No. 69278-3) Lane 2 5 µl PCR Reaction

USA and Canada Tel (800) 526-7319 novatech@novagen.com Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk All Other Countries Contact Your Local Distributor www.novagen.com novatech@novagen.com

2

1

#### Amplification of 919 bp ORF from plasmid DNA

**Reaction setup** 

Component	Volume	Final Concentration
10X Buffer for KOD Hot Start DNA Polymerase	5 µl	1X
25 mM MgSO <sub>4</sub>	3 µl	1.5 mM
dNTPs (2 mM each)	5 µl	0.2 mM (each)
PCR Grade Water	32 µl	
Sense (5') Primer (10 µM)	1.5 µl	0.3 μΜ
Anti-Sense (3') Primer (10 µM)	1.5 µl	0.3 μΜ
Plasmid DNA (10 ng/µl, diluted in TE)	1 µl	0.22 ng/µl
KOD Hot Start DNA Polymerase (1 U/µl)	1 µl	0.02U/µl
Total reaction volume	50 µl	

#### Cycling conditions

Step	Temperature and time
1. Polymerase activation	95°C for 2 min
2. Denature	95°C for 20 s
3. Annealing	55°C for 10 s
4. Extension	70°C for 15 s
Repeat steps 2–4	25 cycles
5. Hold	4°C





1.4 % TAE agarose gel Lane 1 PCR Markers, 50–2000 bp (Cat. No. 69278-3) Lane 2 5 µl PCR Reaction

USA and Canada Tel (800) 526-7319 novatech@novagen.com Germany Tel 0800 100 3496 techservice@merckbiosciences.de United Kingdom and Ireland UK Freephone 0800 622935 Ireland Toll Free 1800 409445 customer.service@merckbiosciences.co.uk